



## Section : 5. Seamless platform

### Module : 5.2. Realtime data preparation and access

#### **i** Understanding where the data in Realtime comes from and how it can be accessed

*“We will review how exposure, aggregation and context data and report templates are sourced and prepared for Realtime and see where to access or maintain them”*

Earthquake, Flood and Volcanic Ash are the hazards currently supported by InaSAFE Realtime. By now you know the origin of the the hazard layers used in InaSAFE Realtime analyses.

But to run an InaSAFE impact function we need exposure layers and optionally, aggregation layers. And to generated the maps in the reports we also need some contextual layers. So where do all these layers come from, since we as users are not selecting them like we do in InaSAFE desktop.

These layers are all carefully prepared and stored on the Realtime server. Then they are programmatically added to each headless InaSAFE Realtime analysis and report.

In the previous version of Realtime they were packaged with the software and 'baked in' and it was difficult to change them. In the new version of Realtime they can be modified much more easily, as long as certain conventions are obeyed, such as file names, geometry types and field names.

The table below shows the layers currently in use and in which modules they are used and what role they play in each module.

You do not have access to change these layers but it is important that you know where they come from and that they can be changed. The layers will however be published on BNPB GeoNode, which will be covered tomorrow.

Data	Type	File Name	Usage in Realtime			Used Field Name	Description
			EQ	Flood	Volcanic Ash		
Place (cities and airports)	Vector point	idn_places_wgs84.shp	Analysis & Context		Analysis & Context	Population	Number of people
						type	Identify which one is airport and which one is place
						name	name of places and airports
Population	Raster	idn_population_200m_wgs84.tif	Analysis & Context		Analysis	-	number of people
Landcover	Vector polygon	idn_landcover_250k_wgs84.shp			Analysis	Classes	Landcover classes
Jakarta Population	Vector polygon	dki_jakarta_population_wgs84.shp		Analysis		Total	Number of people
Volcano	Vector point	idn_volcano_wgs84.shp			Context	Name	Name of volcano
						TimeZone	The region timezone of the volcano crater
						UTC	The UTC timezone of the volcano crater
						Subregion	A region of the volcano crater
						Elevation	The elevation of the volcano crater above sea level
						Province	Province of the volcano crater
Indonesia Administration Boundaries	Vector polyline	idn_admin_boundaries_wgs84.shp	Context	Context	Context	Keterangan	Administrative boundaries
Hillshade	Raster	idn_hillshade_wgs84.tif			Context	-	Hillshade

#### **You try:**

**Goal: In this module we aim to understand which other layers Realtime uses and how it uses them**

Look at an example of each type of report from Realtime: Earthquake, flood and volcanic ash and write down where each of the layers in the table above is used in that report. Look in the maps, tables and other report elements.

Can you find where each one is used?

Compare and discuss with your neighbour.

### **More about Realtime data and reports**

Each layer used in Realtime is described by a metadata document such as the one to the right, for the Land cover layer. The metadata describes the provenance and processing history of the data set. Most of the data layers were not simply downloaded from somewhere and used directly. As with Land cover, many of them required skilled and laborious GIS work to prepare.

Another variable component of realtime is the report templates. These are normal QGIS qpt files with layouts and expressions just like those you've been learning about in InaSAFE desktop. So if you notice an error in a report or want to add some more information, for example, you can modify an existing template and replace it. A lot of care needs to be taken to test it before replacing an existing template. As with the various GIS datasets, you do not have access yet to changing the templates but there should be a web interface in Realtime to upload new templates. Only authorised users will be able to do so.

## Landcover

<b>File Name :</b>	idn_landcover_250k_wgs84.shp
<b>Geometry :</b>	Vector Polygon
<b>Data Type :</b>	Classified
<b>Unit :</b>	Class
<b>Source :</b>	BIG
<b>Source URL:</b>	<a href="http://tanahair.indonesia.go.id/">http://tanahair.indonesia.go.id/</a>
<b>Source Date :</b>	2006
<b>Obtain Date :</b>	2015
<b>Processing Date :</b>	2016
<b>Coverage :</b>	Indonesia
<b>Description :</b>	National landcover dataset compiled by DMInnovation from separate datasets for different landcover types; classified into six classes for standard reporting: <ol style="list-style-type: none"><li>1. <b>Forest</b> consist of <i>Hutan Rimba</i></li><li>2. <b>Plantation</b> consist of <i>Kebun</i> and <i>Ladang</i>.</li><li>3. <b>Rice Field</b> consist of <i>Sawah</i>.</li><li>4. <b>Settlement</b> consist of <i>Permukiman</i>.</li><li>5. <b>Water Supply</b> consist of <i>Danau, Sungai, Tambak, Empang, Waduk, Perairan lainnya</i>.</li><li>6. <b>Other</b> consist of <i>pasir, and semak belukar</i>.</li></ol>

**Field Used :** *Classes* (landcover classes)

**Usage :** These data are used as exposure data for InaSAFE Realtime impact analysis on landcover.

### Check your knowledge:

1. Which analysis uses Places point data?:
  - a. *earthquake*
  - b. *flood*
  - c. *volcanic ash*
2. How many different layers had to be combined to create one national land cover layer?:
  - a. *none, it was acquired as a complete data set*
  - b. *six*
  - c. *three*